Example: A light beam incident on the top surface of a block of plastic $(n_P=1.49)$ makes an angle of 50° with respect to normal. Determine the angle the light makes with respect to the normal as it exits the side of the block into water $(n_W=1.33)$.	Example: A pencil placed 30cm from a spherical mirror yields a real image 6cm from the mirror. (a) Is the mirror concave or convex? (b) Is the image erect or inverted? (c) Determine the radius of curvature of the mirror. (d) Determine the magnification.
Example: A pencil placed 30cm from a spherical mirror yields a real image 6cm from the mirror. (a) Is the mirror concave or convex? (b) Is the image erect or inverted? (c) Determine the radius of curvature of the mirror. (d) Determine the magnification.	Example: An image of a 20.0cm tall object is to be projected on a screen 5.00m from the object. It is important that the image be 2.50m tall. Determine the location, focal length and type of thin lens to use.
Example: A student wants to create a interference pattern on a screen with bright spots separated by a distance of 25cm. She uses a 580nm laser normally incident on a pair of slits separated by a distance of 0.02mm. How far in front of the screen should she place the slits?	Example: A diffraction grating has $800 lines/mm$. A pair of lasers with wavelengths of $\lambda_A = 594.7 nm$ and $\lambda_B = 594.1 nm$ are incident upon a diffraction grating with $800 lines/mm$. How many lines must be illuminated to resolve the lasers in first order?



